

Studies on the nature of the pathogenicity of  
soil-borne Cylindrocarpon species

by

*Mark William*

M.W. Sweetingham, B.Agr.Sc.(Hons.)


Submitted in partial fulfilment of the requirements for the  
Degree of Doctor of Philosophy

University of Tasmania

Hobart

July 1983

This thesis contains no material which has been accepted for any other degree or diploma in any university, and to the best of my knowledge, contains no copy or paraphrase of material previously published or written by any other person, except where due reference is made in the text of the thesis.

A handwritten signature in black ink, appearing to read 'M.W. Sweetingham', with a large, sweeping flourish at the end.

M.W. Sweetingham

University of Tasmania

Hobart

June 1983

# CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENTS	viii
SUMMARY	ix
Abbreviations used throughout the text	xi
1. <u>INTRODUCTION</u>	1
2. <u>LITERATURE SURVEY</u>	3
2.1    Occurrence and saprophytic behaviour of <u>Cylindrocarpon</u> species in soil	3
2.1.1   Isolation from soil	3
2.1.2   Association with the rhizoplane	4
2.1.3   Competitive saprophytic ability	5
2.2    Association of soil-borne <u>Cylindrocarpon</u> species with plant disease	6
2.2.1   Root rots	6
2.2.2   Vascular disorders	9
2.2.3   Damping-off of seedlings	10
2.2.4   Soft rots	11
2.2.5   Stunting and reduced vigour of plants	11
2.3    Studies on the pathogenicity of <u>Cylindrocarpon</u> species	13
2.3.1   Environmental factors influencing pathogenicity	13
2.3.2   The variable virulence of <u>Cylindrocarpon</u> species, and isolates within a species	14
2.3.3   Host specificity of <u>Cylindrocarpon</u> isolates	15
2.3.4   Variable host resistance to <u>Cylindrocarpon</u>	16
2.4    Mechanisms of the pathogenicity of <u>Cylindrocarpon</u>	17
2.4.1   Toxins	17
2.4.2   Enzymes	19
2.4.3   Indole acetic acid	20

	<u>Page</u>
3. <u>EXPERIMENTAL</u>	21
Procedures used throughout the study	21
3.1   Occurrence, identification, and properties of Tasmanian soil-borne <u>Cylindrocarpon</u> species	23
3.1.1 Isolation and identification	23
3.1.1.1 Development of a selective medium	23
3.1.1.2 Recovery from soil and plant tissue	27
3.1.1.3 Description and taxonomy of the isolates	30
3.1.2 Toxin production <u>in vitro</u>	35
3.1.2.1 Rapid qualitative assay for brefeldin A and radicicol	35
3.1.2.2 Isolation of pure brefeldin A and radicicol	39
3.1.3 Pectic enzyme production <u>in vitro</u>	41
3.1.4 The effect of temperature on growth <u>in vitro</u>	47
3.2   Properties of brefeldin A and its possible involvement with plant disease caused by <u>Cylindrocarpon</u> species	51
3.2.1 Antifungal properties	51
3.2.2 Preliminary investigations of phytotoxicity	54
3.2.2.1 Seed germination tests	54
3.2.2.2 Excised leaves	57
3.2.2.3 Tomato plants in solution culture	58
3.2.3 Uptake and translocation of brefeldin A by strawberry plants	63
3.2.4 Stunting of strawberry plants	69
3.2.4.1 Dosage response of strawberry plants to brefeldin A	70
3.2.4.2 Inoculation of strawberry plants with brefeldin A producing isolates of <u>Cylindrocarpon</u>	73

	<u>Page</u>
3.3 A grapevine disease in Tasmania associated with <u>Cylindrocarpon destructans</u> (Zins.) Scholten	85
3.3.1 Disease description	85
3.3.1.1 Introduction	85
3.3.1.2 Isolation of <u>Cylindrocarpon</u> and microscopic examination of diseased tissue	86
3.3.2 Factors affecting infection	90
3.3.2.1 Preliminary inoculation experiments	90
3.3.2.2 Association of <u>Cylindrocarpon</u> with vine roots in southern Tasmania	91
3.3.2.3 Waterlogging stress	92
3.3.3 The role of brefeldin A	96
3.3.3.1 Attempted recovery of brefeldin A from diseased tissue	97
3.3.3.2 Response of vines grown in solution culture amended with brefeldin A	98
3.4 Pathogenicity of <u>Cylindrocarpon</u> species towards <u>Pinus radiata</u> D. Don seedlings	101
3.4.1 Pathogenicity tests in sterile media	101
3.4.2 Pathogenicity tests under non sterile conditions	107
3.5 Macerative pathogenicity	112
3.5.1 Pathogenicity of <u>Cylindrocarpon</u> towards carrot and parsnip roots	112
3.5.1.1 Effect of wounding and inoculum potential	112
3.5.1.2 Variable virulence of <u>Cylindrocarpon</u> isolates	116

	<u>Page</u>
3.5.2 Association of pectic enzymes with tissue maceration	118
3.5.2.1 Factors affecting pectic enzyme production by <u>Cylindrocarpon</u> species <u>in vitro</u>	118
3.5.2.2 Production of pectic enzymes by <u>Cylindrocarpon</u> species in neutral to alkaline media	125
3.5.2.3 Detection of pectic enzymes in diseased tissue	130
4. <u>GENERAL DISCUSSION</u>	137
4.1 Brefeldin A and toxigenic pathogenicity	137
4.1.1 Stunting of seedling growth	137
4.1.2 Reduced vigour of strawberry plants	138
4.1.3 Vascular dysfunction	140
4.2 Pectic enzymes and macerative pathogenicity	140
4.3 <u>Cylindrocarpon</u> as a 'weak' or 'minor' pathogen	142
4.3.1 Environmental factors influencing pathogenicity	143
4.3.2 Disease complexes	144
4.4 Conclusions	145
<u>BIBLIOGRAPHY</u>	147
<u>APPENDICES</u>	169
1.0 Genera of fungi reported to produce brefeldin A	169
1.1 Czapek Dox medium	170
1.2 A list of the <u>Cylindrocarpon</u> isolates studied and their origin	171
1.2b Map of Tasmania showing the locations where <u>Cylindrocarpon</u> was isolated	175

	<u>Page</u>
1.3 Cultural characteristics of the <u>Cylindrocarpon</u> isolates studied	176
1.4 p-Anisaldehyde reagent	179
1.5 Mass spectra of brefeldin A and radicicol	180
1.6 Pectin-acrylamide gel mixture	182
1.7 Amylase zymograms	183
1.8 Ribonuclease zymograms	185
2.1 Antifungal properties of brefeldin A	187
2.2 Standard curve of Flame Ionization Detector response to brefeldin A	190
2.3a Gas chromatogram of the extract from laminas of strawberry plants treated with brefeldin A	191
2.3b Quantitative gas chromatography analysis of brefeldin A in the tissues of strawberry plants treated with the substance	192
2.4a Gas chromatogram of the extract from shoot tissue of strawberry plants inoculated with <u>C. destructans</u>	194
2.4b Quantitative gas chromatography analysis of brefeldin A in shoot tissue of strawberry plants inoculated with <u>C. destructans</u>	195
2.5 Quantitative gas chromatography analysis of brefeldin A in <u>C. didymum</u> inoculum cultured on autoclaved strawberry roots	196
3.1 Spurr's ultra-low viscosity plastic embedding medium	197
3.2 Sensitivity of the GC-FID assay for brefeldin A extracted from diseased grapevine tissue	198
5.1 TBA assay for pectin lyase reaction products: UV absorption spectrum of reaction mixture	199

	<u>Page</u>
5.2 Determination of relative pectin lyase activity of <u>Cylindrocarpon</u> species cultured on NiC medium	200



**ACKNOWLEDGEMENTS**

I thank my supervisor, Professor G.C. Wade, for his encouragement and overall guidance throughout the course of this project.

I would also like to thank members of the technical and professional staff and fellow postgraduate students in the Department of Agricultural Science for their advice and constructive criticism during the course of the project.

I am particularly indebted to Mr. R.H. Cruickshank for introducing me to the techniques of enzyme electrophoresis, Mr. G.W. Leggett for a practical introduction to gas chromatography, and Mr. N. Davies for the operation of the G.C.M.S.

The advice and interest shown by Dr. H.J. Elliott and Dr. C. Palzer of the Tasmanian Forestry Commission was also much appreciated.

Also, I wish to thank Mrs. Sally Jones and Miss Pippa Davis for their assistance with the presentation of the thesis.

Finally, I thank my wife, Sally, for her endless patience, encouragement and support.

## SUMMARY

**SUMMARY**

A study was made of the association of Cylindrocarpon species with the below-ground portions of a range of plants in Tasmania, and of the pathogenic effects of members of the genus.

A selective medium was developed to enhance the recovery of Cylindrocarpon species from soil and plant tissue. Several species were isolated from the root surfaces of a range of apparently healthy woody and herbaceous plants. Cylindrocarpon species were also isolated from the discoloured vascular tissue in the basal stem portion of several diseased woody plant species. C. destructans was the most frequently isolated species, but C. didymum, C. obtusisporum, C. tenue, C. theobromicola, C. candidum and an undescribed species were also recorded. The cultural characteristics and morphology of the isolates were described.

Most of the Cylindrocarpon isolates produced either the metabolite brefeldin A or radicicol when cultured on Czapek Dox medium. The non host-specific phytotoxicity of brefeldin A was confirmed and the spectrum of its antifungal activity was also examined.

A brefeldin A producing strain of C. destructans was consistently isolated from the discoloured wood in the base of the trunks of mature grapevines suffering from a dieback disease. The xylem vessels and functional phloem became blocked with tyloses and gum and fungal hyphae were visible in the ray cells of the diseased tissue. No other potential pathogens were isolated from the diseased tissue. Field observations suggested that the waterlogging of the trunks at ground-

level was necessary for the disease to develop. A waterlogging stress was required for the fungus to reproduce the characteristic disease symptoms on inoculated six-month-old vine cuttings. However, brefeldin A was not detected in naturally diseased grapevine tissue and could not be demonstrated to have a role in this disorder.

Certain brefeldin A producing strains of Cylindrocarpon were capable of parasitizing the root cortex of radiata pine seedlings under sterile conditions, causing a suppression in root growth. Radiata pine seed germinated in the presence of brefeldin A similarly produced stunted roots. Likewise, the growth of strawberry plants was significantly reduced, in the absence of other disease symptoms, when grown in sterilised soil amended with a brefeldin A producing isolate of C. destructans. This effect was reproduced by growing strawberry plants in soil amended with brefeldin A. Brefeldin A was absorbed by the strawberry roots and translocated into the shoots.

All the Cylindrocarpon isolates were shown to produce a spectrum of pectin degrading isoenzymes when cultured on appropriate media. Inoculation experiments showed several Cylindrocarpon species were capable of producing "soft rot" type lesions on carrot roots. Several species could also infect the hypocotyl of radiata pine seedlings in sterile media, causing post emergence damping-off. Pectin lyase enzymes were detected in the lesions from both carrot roots and radiata pine hypocotyls and may be responsible for the tissue maceration. However, virulence of the isolates was not related to pectin lyase production in vitro.

## ABBREVIATIONS USED THROUGHOUT THE TEXT

### CHEMICALS AND MEDIA:

Tris:	tris(hydroxymethyl)aminomethane
PVP(40):	polyvinylpolypyrrolidone (Average molecular weight: 40,000)
PDA:	potato dextrose agar
CDA:	Czapek-Dox agar

### UNITS:

m:	metre	kg:	kilogram
cm:	centimetre	g:	gram
mm:	millimetre	mg:	milligram
$\mu$ :	micron ( $=10^{-6}$ metre)	$\mu$ g:	microgram
nm:	nanometer	ng:	nanogram
		M:	Molar concentration (moles per litre)
L:	litre	ppm:	parts per million (concentration)
ml:	millilitre	s:	second
$\mu$ l:	microlitre	$^{\circ}$ C:	degrees Centigrade
		$\mu$ E:	microeinstein
		Amp f.s.d.:	Ampere (full scale deflection)
		kV:	kilovolt
		eV:	electron volt

### STATISTICS

S.E. of mean:	standard error of the mean
LSD ( $p = 0.05$ ):	least significant difference at the 5% probability level
LSD ( $p = 0.01$ ):	least significant difference at the 1% probability level